

# IONIC LIQUIDS AS REPLACEMENTS FOR N-DODECANE: A COMPUTATIONAL STUDY OF SOLVENT EFFICACY IN THE PUREX PROCESS

<sup>1</sup>Chukwuemeka Nkemdilim Obi and <sup>2</sup>Ibrahim Yusuf Ahmed

## Article Info

**Keywords:** Ionic Liquids, PUREX Process, Solvent Replacement, Distribution Coefficient, Hydrophobicity

## DOI

10.5281/zenodo.12806252

## Abstract

This study investigates the potential of ionic liquids as alternative solvents to the volatile n-dodecane used in the PUREX process for separating uranium and plutonium from spent nuclear fuel. Using the COSMO-RS method, 2106 ionic liquids were screened to evaluate their suitability. Key parameters, including the distribution coefficients of uranyl nitrate in an ionic liquid/aqueous nitric acid biphasic system and the solubility of these ionic liquids in water, were assessed. Seventeen ionic liquids, particularly those based on tetrafluoroborate (BF<sub>4</sub>), were identified as highly hydrophobic. Among these, hexamethyl guanidium tetrafluoroborate demonstrated the highest distribution coefficient, surpassing that of conventional n-dodecane despite its increased hydrophilicity. The findings indicate that certain ionic liquids hold promise as effective replacements for n-dodecane in the PUREX process, potentially enhancing solvent performance in nuclear fuel reprocessing.

## INTRODUCTION

The products contained in SNF are radioactive with halflives from days to millions of years which give a lot of threat to human health and environment (Terry, 2008). Uranium and Plutonium are conventionally recovered through PUREX process. PUREX, an abbreviation for Plutonium (P) Uranium (UR) Extraction (EX), makes use of about 30% tri-*n*-butyl phosphate (TBP) with Kerosene or pure *n*-Dodecane as diluents (Sung et al., 2010). In this process, Uranium and Plutonium are extracted as UO<sub>2</sub><sup>2+</sup> and Pu<sup>4+</sup> and then converted to UO<sub>2</sub> and PuO<sub>2</sub>. Aside from uranium and plutonium, various fission products can be recovered from SNF. Thus some extraction processes have been developed using different extracting agents to recover fission products from PUREX raffinate. The extraction processes developed for SNF treatment and recovery, utilize organic solvents that are volatile, hazardous, toxic, flammable, and unstable towards the radiation and heat generated by the SNF (Sung et al., 2010). Since the undesirable properties of these conventional solvents may pose several threats and problems

<sup>1</sup> Nigerian National Petroleum Cooperation (NNPC), Nigeria.

<sup>2</sup> Chemical Engineering Department, Ahmadu Bello University Zaria.

both to plant operation and to environment, several researches are ongoing in search of alternate safer solvents such as ionic liquids (ILs) which have unique physical and chemical properties, such as specific solvent abilities, negligible vapor pressures, non-flammability, thermal and radiological stability and broad liquid temperature ranges among many. They have led to promising applications as environmentally benign solvents that can be possible candidates to replace conventional organic solvents (Sung et al., 2010).

During the past 10 years, ionic liquids (ILs) have received increased attention. They are organic salts, and their chemical and physical properties can be tailored by the selection of an anion and cation (Wasserscheid, 2000). Therefore, it is possible to generate a huge number of different ionic liquids, each with specific properties. Despite the interest, accurate thermodynamic data of ionic liquids and their mixtures are still rare and to exploit the potential of these new substances, it would be of great value to have prediction methods that can reliably predict the thermodynamic properties of ionic liquids and their mixtures. This would help to scan the growing set of already known ILs in order to find suitable candidates for a certain task or to design new ILs for special applications (Geetanjali and Kumar, 2008). Group contribution methods, which are the most widely used theoretical models are not applicable because group interactions parameters are not available at present for ionic liquids and the group contribution concept is not suitable to handle the long-range interactions in ionic compounds (Hamad and Sumon, 2008). Monte Carlo simulations and molecular dynamics need appropriate force-fields for the treatment of ionic liquids, which have to be developed (Hamad and Sumon, 2008). This work will present the application of a continuum solvation based thermodynamic prediction model, Conductor-like Screening Model for Real Solvents, COSMO-RS for prediction of thermo-physical data which is a computational approach independent of experimental data and with general applicability (Eckert and Klamt, 2002).

The aim of the research is to computationally screen suitable hydrophobic ionic liquids as solvents for the extraction of Uranium from spent nuclear fuel using COSMO –RS method by determining the distribution coefficients of uranyl nitrate in hydrophobic ionic liquids/ aqueous nitric acid biphasic solution and determination the solubility of the ionic liquids in water. The work will be limited to the application of COSMO-RS, a thermodynamic model which uses statistical thermodynamic approach in determination of thermodynamic properties based on results obtained from quantum chemical/COSMO computation for the screening of ionic liquids for Uranium Extraction from spent nuclear fuel.

## **MATERIALS AND METHODS**

The following are the materials used in carrying out the research work:

- a) System Softwares
  - i) Linux operating system
  - ii) Windows 7 operating system
- b) Application Softwares
  - i) Chem draw 3D ultra 8.0/ Marvin sketch software. ii) TURBOMOLE 5.6 Program Package (LINUX VERSION). iii) COSMO therm Software.

The following are the methods used in carrying out the research work:

- 1) The Marvin sketch software is first used in sketching the structure of each individual molecule. The environment has a drawing kit including all elements in the periodic table and heterocyclic structures for easy sketch of the molecular structures. Marvin sketch software operates on windows. The file generated is saved in xyz format and is used as an input file for TURBOMOLE to perform the quantum chemical/COSMO calculation.
- 2) The TURBOMOLE uses the xyz file generated from Marvin sketch to perform quantum chemical/COSMO calculation to generate the screening charge densities of the molecules. The screening charge is a microscopic property of a molecule like internal energy and it is the main parameter used in the COSMORS

model. The quantum chemical/COSMO computation on TURBOMOLE 5.6 is parameterized using DFT-level of computation utilizing BP-functional on TZVP basis set. The file is saved as a COSMO file and an input file to COSMO therm to perform the statistical thermodynamics and to determine the screening charge density profiles and chemical potentials of the individual molecules (Klamt, 1995). The higher the specification of the computer machine, the faster the computation of the screening charges of the molecules.

- 3) A database is created for all the COSMO files generated using the TURBOMOLE program software.
- 4) The screening charge density profiles and the chemical potentials of the molecules are determined using COSMO therm. The COSMO therm is the software that solves the COSMO-RS model based on self consistency field algorithm (SCF) to statistically determine the screening charge density profiles and the chemical potentials of the molecules (Klamt, 1995).
- 5) The 81 cations and 26 anions were combined to generate 2106 ionic liquids (Table 1).
- 6) The solubility of the ionic liquids in water is determined from COSMO therm using the equation;

$$\text{Log}_{10}(x_j^{\text{sol}}) = [\mu_j^{(\text{p})} - \mu_j^{(\text{l})} - \max(0, \Delta G_{\text{fus}})] / (RT \ln(10))$$

Where  $\Delta G_{\text{fus}}$  is the heat of fusion per mole. If the compound is solid, the energy change of a compound from the

S/N	Anions		super cooled liquid state
1	Tetrafluoroborate	14	bissalicylatoborate to the
2	Hexafluorophosphate (vi)	15	tetracyanoborate ordered
3	methyl sulfate	16	bis(2,4,4-trimethylpentyl)phosphinate solid state
4	chlorate	17	bis-pentafluoroethyl-phosphinate has to be
5	bromate	18	bis(trifluoromethylsulfonyl)methane taken into
6	Iodide	19	decanoate account, that
7	formate	20	tosylate is, the

solutes

Gibbs free energy of fusion  $\Delta G_{\text{fus}}$  (or equivalently its Gibbs free energy of crystallization  $\Delta G_{\text{cryst}} = -\Delta G_{\text{fus}}$ ) has to be either given and computed from experimental data or estimated by COSMOtherm. For a liquid, it is automatically specified but for solids it has to be determined using QSPR and then specified manually.

$T$  = Temperature in degree Celsius  $R$  = molar gas constant in KJ/mol/Kelvin  $\mu_j^{(\text{i})}$  = chemical potential of solute  $j$  in solvent  $i$  in energy/mole

$\mu_j^{(\text{p})}$  = chemical potential of pure compound  $j$  in energy/mole  $x_j$  = mole fraction of the solubility of solutes in water (dimensionless).

- 7) The distribution coefficient and octanol-water partition coefficients are determined from COSMOtherm using the equation:

$$\text{Log}_{10}(P) = \text{Log}_{10}[\exp(\mu_j^{(1)} - \mu_j^{(2)}) / RT] \cdot V_1/V_2$$

Where  $v_1$  and  $v_2$  are volume quotients of phase 1 and phase 2  $\mu_j^{(1)}$  = chemical potential of uranyl nitrate in phase 1 in energy/mole  $\mu_j^{(2)}$  = chemical potentials of uranyl nitrate in phase 2 in energy/mole  $P$  = partition coefficient (dimensionless).

**Table 1.** List of cations and anions.

S/N	Cations
-----	---------

1	1,1-dimethyl-pyrrolidinium	42	methyl-trioctyl-ammonium
2	1,1-dipropyl-pyrrolidinium	43	tetra-methyl ammonium
3	1-ethyl-1-methyl-pyrrolidinium	44	tetra-ethyl ammonium
4	1-butyl-1-methyl-pyrrolidinium	45	tetra-n-butyl ammonium
5	1-butyl-1-ethyl-pyrrolidinium	46	benzyl-triphenyl-phosphonium
6	1-hexyl-1-methyl-pyrrolidinium	47	tetrabutyl-phosphonium
7	1-octyl-1-methyl-pyrrolidinium	48	triethyl-tetradecyl-phosphonium
8	3-methyl-imidazolium	49	triisobutyl-methyl-phosphonium
9	1-butyl-imidazolium	50	guanidinium
10	1,3-dimethyl-imidazolium	51	hexamethylguanidinium
11	1-ethyl-3-methyl-imidazolium	52	N,N,N,N,N-pentamethyl-N-isopropyl-guanidinium
12	1-butyl-3-methyl-imidazolium	53	N,N,N,N,N-pentamethyl-N-propyl-guanidinium
13	1-pentyl-3-methyl-imidazolium	54	N,N,N,N-tetramethyl-N-ethylguanidinium
14	1-hexyl-3-methyl-imidazolium	55	S-ethyl-N,N,N,N-tetramethylisothiouronium
15	1-octyl-3-methyl-imidazolium	56	O-ethyl-N,N,N,N-tetramethylisouronium
16	1-decyl-3-methyl-imidazolium	57	O-methyl-N,N,N,N-tetramethylisouronium
17	1-dodecyl-3-methyl-imidazolium	58	N-butyl-isoquinolinium
18	1-tetradecyl-3-methyl-imidazolium	59	morpholinium
19	1-hexadecyl-3-methyl-imidazolium	60	methylmorpholinium
20	1-octadecyl-3-methyl-imidazolium	61	dimethylmorpholinium
21	1-benzyl-3-methyl-imidazolium	62	N-fluoropropoxy-methylmorpholinium
22	1-ethyl-2-3-methyl-imidazolium	63	O-ethyl-tetrapropylisouronium
23	1-propyl-2-3-methyl-imidazolium	64	O-hydro-tetraethylisouronium
24	1-butyl-2-3-methyl-imidazolium	65	O-hydro-tetramethylisouronium
25	1-hexyl-2-3-methyl-imidazolium	66	O-methyl-tetraethylisouronium
26	1-hexadecyl-2-3-methyl-imidazolium	67	O-methyl-tetrapropylisouronium
27	1-methyl-3-(3-phenyl-propyl)-imidazolium	68	O-propyl-tetramethylisouronium
28	4-methyl-n-butylpyridinium	69	O-propyl-tetrapropylisouronium
29	1-ethyl-pyridinium	70	O-butyl-tetramethylisouronium
30	1-butyl-pyridinium	71	O-ethyl-tetraethylisouronium
31	1-hexyl-pyridinium	72	diethyl-dimethylammonium
32	1-octyl-pyridinium	73	dimethylammonium
33	1-butyl-3-ethyl-pyridinium	74	ethyl-trimethylammonium
34	1-butyl-3-methyl-pyridinium	75	methyl-triethylammonium
35	1-butyl-4-methyl-pyridinium	76	tetra-ethyl ammonium
36	1-hexyl-3-methyl-pyridinium	77	tetramethylammonium

37	1-hexyl-4-methyl-pyridinium	78	trimethylammonium
38	3-methyl-1-octyl-pyridinium	79	n-methyldiazabicyclo-undec-7-enium
39	4-methyl-1-octyl-pyridinium	80	n-ethyldiazabicyclo-undec-7-enium
40	1-butyl-3,4-dimethyl-pyridinium	81	n-hexyldiazabicyclo-undec-7-enium
41	1-butyl-3,5-dimethyl-pyridinium		

**Table 1.** Contd.

8	toluene-4-sulfonate	21	bis(pentafluoroethylsulfonyl)imide
9	trifluoromethane-sulfonate	22	n-methyl-n-butylcarbamate
10	tris(nonafluorobutyl)trifluorophosphate	23	n-methyl-n-propylcarbamate
11	tris(pentafluoroethyl)trifluorophosphate	24	Nitrate
12	bisbiphenyldiolatoborate	25	Thiocyanate
13	bisoxalatoborate	26	Methylphosphonate

**Post processing**

From the properties determined, criteria for the selection of suitable ionic liquids were imposed. These are based on the following criteria;

***Hydrophobicity***

Out of the 2106 ionic liquids, those with mole fraction solubility of 1 are considered hydrophilic while those with mole fraction solubility of zero are considered as hydrophobic ionic liquids.

***Comparable partition coefficient with n-dodecane***

Those ionic liquids with distribution coefficient greater than or equal to that of n-dodecane were recommended for screening. Similarly, those with solubility mole fraction of less than 1 are considered partially miscible with water while those with solubility mole fraction of 1 are considered fully miscible with water (hydrophilic). Those ionic liquids that satisfy criteria 1 and 2 were recommended. Using these criteria, 111 ionic liquids were screened out of the in the database. 17 ionic liquids based on BF<sub>4</sub> are the most hydrophobic.

The 17 ionic liquids are given below:

- 1) 1,1-dimethyl-pyrrolidinium tetrafluoroborate
- 2) 1-ethyl-1-methyl-pyrrolidinium tetrafluoroborate
- 3) tetra-ethylammoniumtetrafluoroborate
- 4) hexamethylguanidiniumtetrafluoroborate
- 5) N,N,N,N,N-pentamethyl-N-isopropyl-guanidiniumtetrafluoroborate
- 6) N,N,N,N,N-pentamethyl-N-propyl-guanidiniumtetrafluoroborate
- 7) S-ethyl-N,N,N,N-tetramethylisouroniumtetrafluoroborate
- 8) O-ethyl-N,N,N,N-tetramethylisouroniumtetrafluoroborate
- 9) O-methyl-N,N,N,N-tetramethylisouroniumtetrafluoroborate
- 10) O-methyl-tetraethylisouroniumtetrafluoroborate
- 11) O-propyl-tetramethylisouroniumtetrafluoroborate
- 12) O-butyl-tetramethylisouroniumtetrafluoroborate
- 13) O-ethyl-tetraethylisouroniumtetrafluoroborate

- 14) methyl-triethylammoniumtetrafluoroborate
- 15) n-methyldiazabicyclo-undec-7-eniumtetrafluoroborate
- 16) n-ethyldiazabicyclo-undec-7-eniumtetrafluoroborate
- 17) tetra-methylammoniumtetrafluoroborate

## RESULTS AND DISCUSSION

### Data validation of solubility


The experimental solubility in water values of some solvents determined was compared with that of COSMORS method. The correlation factor obtained was 0.994 which shows a good correlation between the experimental and that of the model. This is a validation of the model for determination of solubility of compounds in water. Figure 1 gives the comparison between the experimental and the model results. The experimental results were obtained from Laurie (2004) and Felicia et al. (2009).

### Evaluation of the distribution coefficient and solubility of n-dodecane and the screened ILs


The distribution coefficient of uranyl nitrate in n-dodecane /aqueous nitric acid biphasic system and that of uranyl nitrate in ionic liquids/aqueous nitric acid biphasic system was determined using COSMO-RS method. The solubility of the n-dodecane and ionic liquids in water was also determined to determine the hydrophobicity of the ionic liquids and the n-dodecane. As shown in Figure 2, the solubility of n-dodecane in water is 0.0072 and is the most hydrophobic among the solvents and this gives its advantage of easier recycle and recovery. This is because the more hydrophobic the solvent is, the better the separation with the aqueous phase. Among the ionic liquids screened, 17 of them are the most hydrophobic closest to n-dodecane with solubility mole fraction of less than 0.1. Although they have comparable or higher distribution coefficient with the conventional solvent, n-dodecane. The advantage of higher distribution coefficient is the use of less number of stages during continuous extraction. It can also be seen that some ionic liquids give high distribution coefficient but more miscible with water. It has been reported that hydrophilic ionic liquids forms complex with uranyl nitrate via anionic and cationic complexes with contamination of the aqueous phase and this is a limitation in the application of these types of ionic liquids for this application despite the high distribution coefficient (Binnemans, 2007). Hydrophobic ionic liquids on the other hand, forms complex via neutral complexes without contamination of the aqueous phase (Binnemans, 2007).

### Effect of nitric acid concentration

Nitric acid concentration affects the distribution coefficient




**Figure 1.** Comparison of experimental solubility values against COSMO-RS.




**Figure 2.**  $K_d$  (Uranyl nitrate) against the solubility of solvents in water.  
 ◆  $\text{Bf}_4$  based ILs    ▲  $\text{TF}_2\text{N}$  based ILs    ● Thiocyanate based ILs  
 ■ Iodo based ILs    □ pentafluoroethylphosphinate based ILs  
 ▲ Methylsulfate based ILs    ■ n-dodecane (Reference Solvent).

of the uranyl nitrate in the ionic liquid/aqueous nitric acid acid concentration from 0.2 to 4 molar and then slightly biphasic system as can be seen in Figure 3. The decreases to 8 molar. This is because with increase in distribution coefficient increases with increase in nitric nitric acid concentration, the solubility of the solute in the aqueous phase decreases as a result of poor coordination of the nitrate ions with the uranyl nitrate while its solubility in the ionic liquid phase increases. Conversely, the solubility of the solute increases with the increase in nitric acid concentration as a result of its good coordination with the nitrate ions.



**Figure 3.** Effect of nitric acid concentration on the  $D (UO_2NO_3)_2$  J, K of the 17 selected ionic liquids.



**Figure 4.** Effect of TBP concentration on the  $D (UO_2NO_3)_2$  J, K of the selected ionic liquids.

#### Effect of TBP concentration on distribution coefficient

TBP is the universally accepted extractant used in the PUREX process for the extraction of uranium and plutonium from spent nuclear fuel. It forms complex with elements at the +vi and +iv oxidation states. Uranium with the most stable oxidation state of +vi forms complex with the TBP while plutonium at the +iv oxidation state also forms complex with the TBP. They are extracted together to the organic phase and then separated to obtain uranium and plutonium. As shown in Figure 4, the distribution coefficient of uranyl nitrate in ionic liquids/aqueous nitric acid biphasic system, increases with increase in TBP concentration from 20 to 30% and then decreases with increase in TBP concentration from 30 to 45%. The maximum distribution coefficient is obtained at 30% TBP concentration. This is also reported by Binnemans (2007) that optimum distribution coefficient is obtained at 30% TBP concentration from the range of percentage application of the TBP concentration from 20 to 40%.

#### Conclusion

- 1) There are many potential ionic liquids that display higher distribution coefficient than n-dodecane.

- 2) 17 ionic liquids based on  $\text{BF}_4$  are the most recommended hydrophobic ionic liquids of the 111 screened.
- 3) Tributyl-phosphate TBP has effect on the distribution coefficient. The highest distribution coefficient is obtained at 30% TBP by volume and four (4) molar concentration trioxonitric (vi) acid.
- 4) The effect of temperature with the distribution coefficients was also investigated. The distribution coefficients of all the ionic liquids/nitric acid biphasic system were decreasing with temperature from 25 to 100°C.

## REFERENCES

- Binnemans K (2007). Lanthanides and Actinides Chemistry in ionic liquids. Chem.Rev.
- Eckert F, Klamt A (2002). Fast solvent screening via quantum chemistry COSMO-RS approach. AIChE J. 48(2):369.
- Felicia S, Alice M, Petre I, Adrian B, Titus C, Alexandru T B (2009). New alternatives for estimating the Octanol/water partition coefficient and water solubility for volatile organic compounds using GLC data (Kovats retention indices). ARKIVOC (10):174-194.
- Hamad EM, Sunon KZ (2008). Computational screening of ionic liquids for aromatic extraction using COSMO-RS.
- Klamt A (1995). Conductor-like Screening Model for Real Solvents: A New Approach to the Quantitative Calculation of Solvation Phenomena. J. Phys. Chem. 99:2224.
- Sung HH, Russel MN, Yoon MK (2010). Reprocessing of spent nuclear fuel using ionic liquid. Korean J. Chem. Eng. 27(9):1360-1365
- Wasserscheid P, Keim W (2000). Ionic liquids - New solutions for transition metal catalysis. Angew. Chem.-Int. Edit. 39(21):3773-3789.